



# Somerset Surgical Specialists

## Pre-Endovascular Ablation (Closure) Instructions

Day of Procedure:

- Please arrive to the office 15 mins prior to your scheduled procedure time.
- Take 1 (one) Xanax pill one hour prior to schedule procedure. **Bring second pill with you** to the office. You will be directed when it should be taken.
- **Bring compression stockings with you** to the office (thigh high, 20-30 mmHg strength)
- Arrange for someone to accompany you. You will not be able to drive after the procedure.

**All medications will be called into the pharmacy we have on file.**

## Post-Endovascular Ablation (Closure) Instructions

Please be on your feet as much as possible today. We prefer that you do not go home and lie down. To assist in the proper closure of your vein, we want as much pressure on the vein to help keep it closed. Running errands, doing chores around the house, or going for a walk in your neighborhood are some examples of accomplishing this goal.

For the next 72 hours, you will need to keep your compression stocking on at all times with the exception of bathing.

24-hours after your procedure, you may remove your compression stocking to bathe. Once you have completed bathing, the stocking must go back on. Bandages may be removed and they **DO NOT** need to be replaced.

**REMEMBER: the compression stocking must stay on around the clock for the next three days.**

After 72 hours, you will need to wear the compression stocking only during the day and it may be removed before sleep.

Sarah Bryczkowski, MD, FACS

Obi J Imegwu, MD, FAC

Pooja Patel, MD

Thangamani Seenivasan, MD, FACS

30 Rehill Avenue, Suite 3400, Somerville, NJ 08876

(P) 908-725-2400 (F) 908-927-8990



# Somerset Surgical Specialists

Please refrain from any heavy activity such as running and weight lifting for at least three days. Too much pressure against the vein may reopen it and reverse the effect of the procedure.

It is important that you follow up with two ultrasounds and one office visit. Ultrasounds are preferred at RWJ University Hospital-Somerset Vascular Lab. To schedule these appointments please call 1-844-795-4968. \* **SEE ENCLOSED PRESCRIPTIONS. YOU WILL NEED TO BRING THESE WITH YOU TO YOUR APPOINTMENTS\***

- 1st Ultrasound: 3 days from procedure
- 2nd Ultrasound: 10 days from procedure
- Office Visit: 2-3 weeks from your 2nd Ultrasound

During your office visit, the doctor will review your ultrasound reports and determine whether or not the compression stocking will need to be worn for an additional period of time.

If you have any questions, please contact our office at 908-725-2400.

Sarah Bryczkowski, MD, FACS

Pooja Patel, MD

Obi J Imegwu, MD, FAC

Thangamani Seenivasan, MD, FACS

30 Rehill Avenue, Suite 3400, Somerville, NJ 08876

(P) 908-725-2400 (F) 908-927-8990